

WHEN YOUR GOAL IS PARTIAL-ONSET SEIZURE CONTROL, ONCE-A-DAY OXTELLAR XR MAY BE ABLE TO HELP.

Not actual patient.
Used for illustrative purposes.

PAY AS LITTLE AS

\$0 CO-PAY*

ONCE-A-DAY

Oxtellar XR[®]

(oxcarbazepine) extended-release tablets
600 mg 300 mg 150 mg

INDICATION

Oxtellar XR[®] (oxcarbazepine) extended-release tablets is a prescription medicine used to treat partial-onset seizures in adults and children 6 years of age and older.

Please refer to the Important Safety Information (pages 8 and 9) and the Medication Guide for more information on Oxtellar XR, or visit www.OxtellarXR.com/patient.

*For full terms and conditions, please see the Oxtellar XR co-pay savings card, or visit www.OxtellarXR.com.

Epilepsy Facts

1 in 26

Number of Americans who will be diagnosed with epilepsy at some point in their lifetime.

150,000

Number of new epilepsy cases diagnosed in the US each year.

60%

Percentage of people with epilepsy who experience partial-onset seizures.

How are partial-onset seizures different?

Not actual patient.
Used for illustrative purposes.

Partial-onset seizures affect a specific area on one side of the brain. Generalized-onset seizures affect both sides. A majority of people with epilepsy suffer from partial-onset seizures.



The partial-onset seizure medication your healthcare provider prescribes first can make a difference in reducing your seizures or even in becoming seizure-free.

Choosing A Therapy

Finding the right partial-onset seizure medication is important.

Literature describing epilepsy treatments recommends using monotherapy—just 1 seizure medication—in most cases. Increasing the number of seizure medications has its downsides:



SIDE EFFECTS

There is a potential to have more drug interactions and side effects.



COST

It may be more expensive to take additional seizure medications.



MEMORY

It may become harder to remember to take all your seizure medications, which could lead to missed doses.

11 positive clinical trials

Oxtellar XR is the only antiepileptic drug that provides 24-hour controlled delivery of oxcarbazepine, the medicine with powerful evidence for the treatment of partial-onset seizures.



2

Please refer to the **Important Safety Information (pages 8 and 9)** and the **Medication Guide** for more information on Oxtellar XR, or visit www.OxtellarXR.com/patient.


ONCE-A-DAY
Oxtellar XR
(oxcarbazepine) extended-release tablets
600 mg 300 mg 150 mg

3


Extended-Release Designed for 24-Hour Drug Delivery

Solutrol[®] extended-release technology means you take just 1 dose, once a day.

Each tablet of Oxtellar XR contains the drug (oxcarbazepine) and an agent that helps the tablet dissolve evenly and in a controlled manner to provide medication over 24 hours.




Not actual patient.
Used for illustrative purposes.



- Drug
- Solubilizing agent
- Other components

For illustration purposes only; does not represent Oxtellar XR or the actual time medicine is released.

Delivery of 1200 mg once-a-day Oxtellar XR over 24 hours.



Monohydroxy derivative (MHD) becomes the active part of oxcarbazepine once the medicine enters the body.

This study was done in healthy adults. The graph represents sample activities over the course of a day. Oxtellar XR has not been shown to have an impact on these activities.

What would you do between doses of Oxtellar XR?



Simple Once-a-Day Regimen

Taking Oxtellar XR is simple—once a day for full 24-hour medication coverage.

- Your healthcare team can start you on 600 mg/day of Oxtellar XR and increase your dose weekly by 600 mg
- The recommended maintenance dose for once-a-day Oxtellar XR is 1200 mg to 2400 mg once a day

WEEK 1

600 mg/day

600

DAY 1 DAY 2 DAY 3 DAY 4 DAY 5 DAY 6 DAY 7



WEEK 2

1200 mg/day

600


600

DAY 8 DAY 9 DAY 10 DAY 11 DAY 12 DAY 13 DAY 14



Take once-a-day Oxtellar XR on an empty stomach.

For optimal tolerability, take Oxtellar XR at least 1 hour before or at least 2 hours after a meal.



- Take Oxtellar XR once every day
- Take at least 1 hour before or at least 2 hours after a meal
- Do not cut, crush, or chew the tablets before swallowing
- Take Oxtellar XR at the same time each day
- Do not stop taking Oxtellar XR without talking to your healthcare team. Stopping Oxtellar XR suddenly can cause serious problems, including seizures that will not stop (status epilepticus).
- Take Oxtellar XR exactly as prescribed



6

Please refer to the Important Safety Information (pages 8 and 9) and the Medication Guide for more information on Oxtellar XR, or visit www.OxtellarXR.com/patient.

ONCE-A-DAY
Oxtellar XR
(oxcarbazepine) extended-release tablets
600 mg 300 mg 150 mg

7

Important Safety Information

IMPORTANT SAFETY INFORMATION

Do not take Oxtellar XR if you are allergic to oxcarbazepine, or any of the other ingredients in Oxtellar XR, or to eslicarbazepine acetate.

Oxtellar XR can cause serious side effects, including a low level of sodium in your blood. Symptoms of low blood sodium include nausea, tiredness, lack of energy, headache, confusion, and more frequent or more severe seizures. Similar symptoms that are not related to low sodium may occur from taking Oxtellar XR.

The most common side effects include dizziness, sleepiness, headache, balance problems, tremors, vomiting, double vision, and weakness. These are not all of the possible side effects of Oxtellar XR. For more information, ask your healthcare provider or pharmacist.

Like other antiepileptic drugs, Oxtellar XR may cause suicidal thoughts or actions in a very small number of people, about 1 in 500. Before you take Oxtellar XR, tell your healthcare provider if you have or have had depression, mood problems, or suicidal thoughts or behavior. Call a healthcare provider right away if you have thoughts about suicide or dying, attempts to commit suicide, have new or worse depression or new or worse anxiety, feel agitated or restless, experience panic attacks, trouble sleeping (insomnia), or new or worse irritability, feel or act more aggressive, angry, or violent, act on dangerous impulses, have an extreme increase in activity and talking (mania), or experience other unusual changes in behavior or mood.

Oxtellar XR may also cause allergic reactions or serious problems which may affect organs and other parts of your body like the liver or blood cells. You may or may not have a rash with these types of reactions. Call your healthcare provider right away if you have any of the following: swelling of your face, eyes, lips, or tongue, trouble swallowing or breathing, a skin rash, hives, fever, swollen glands, or sore throat that does not go away or comes and goes, painful sores in the mouth or around your eyes, yellowing of your skin or eyes, unusual bruising or bleeding, severe fatigue or weakness, severe muscle pain, frequent infections, or infections that do not go away. Many people who are allergic to carbamazepine are also allergic to Oxtellar XR. Tell your healthcare provider if you are allergic to carbamazepine.

Before taking Oxtellar XR, tell your healthcare provider about all your medical conditions, including if you have or have had suicidal thoughts or actions, have depression or mood problems, have liver problems, have kidney problems, are allergic to carbamazepine, use birth control medicine, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. Oxtellar XR passes into breast milk. Talk with your healthcare provider about the best way to feed your baby if you take Oxtellar XR.

Oxtellar XR may harm your unborn baby. Tell your healthcare provider right away if you become pregnant while taking Oxtellar XR. If you use birth control medicine, Oxtellar XR may cause your birth control medicine to be less effective.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Taking Oxtellar XR with certain other medicines may cause side effects or affect how well they work. Do not start or stop other medicines without talking to your healthcare provider. Especially tell your healthcare provider if you take: carbamazepine, phenobarbital, phenytoin, or birth control medicine.

Do not stop taking Oxtellar XR without talking to your healthcare provider. Stopping Oxtellar XR suddenly can cause serious problems, including seizures that will not stop. Take Oxtellar XR exactly as prescribed.

INDICATION

Oxtellar XR[®] (oxcarbazepine) extended-release tablets is a prescription medicine used to treat partial-onset seizures in adults and children 6 years of age and older.

Please refer to the full [Prescribing Information](#) and [Medication Guide](#) for additional important information.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.



Explore these helpful resources on epilepsy.

You should always talk to your healthcare provider about any questions you may have and for information about the partial-onset seizure treatment plan specifically designed for you. There are also many websites where you can find additional information. Below is a list of resources you may find helpful:

Epilepsy Foundation
Epilepsy.com

Centers for Disease Control and Prevention
CDC.gov/epilepsy

National Association of Epilepsy Centers
NAEC-epilepsy.org

American Epilepsy Society
AESnet.org

The websites listed are not affiliated with, controlled, or endorsed by Supernus Pharmaceuticals, Inc.

Supporting you beyond the prescription.

We're proud to offer Supernus Support, a single point of contact for all of your questions regarding Oxtellar XR, including medication access, insurance coverage, and financial aid for eligible patients. Call Supernus Support to check your availability.

Call us at 1-866-398-0833



All commercially approved patients pay as little as \$0 co-pay.*

Ask your healthcare team for a co-pay savings card, or download one at www.OxtellarXR.com/copay.

Helpful hints to stay on track



Use alarms, smartphone apps, pill boxes, or daily activities such as brushing your teeth as prompts to take your medicine



Continue taking your medicine even when you feel it isn't working or you don't need it



Ask your healthcare provider if there are ways to modify the number of doses you need to take to achieve your treatment goals



*For full terms and conditions, please see the Oxtellar XR co-pay savings card, or visit www.OxtellarXR.com.

10 Please refer to the **Important Safety Information (pages 8 and 9)** and the **Medication Guide** for more information on Oxtellar XR, or visit www.OxtellarXR.com/patient.



11

COULD OXTELLAR XR BE RIGHT FOR YOU?



Not actual patient.
Used for illustrative purposes.

**Ask your healthcare provider about
once-a-day Oxtellar XR and visit
www.OxtellarXR.com/patient to learn more.**

ONCE-A-DAY

Oxtellar XR[®]
(oxcarbazepine) extended-release tablets
600 mg 300 mg 150 mg

**Please refer to the Important Safety Information
(pages 8 and 9) and the [Medication Guide](#)
for more information on Oxtellar XR,
or visit www.OxtellarXR.com/patient.**

Oxtellar XR and Solutrol are registered trademarks of Supernus Pharmaceuticals, Inc. All other trademarks are the property of their respective owners.

© 2021 Supernus Pharmaceuticals, Inc.
All rights reserved. OXT.2021-0059.

 **Supernus**[®]
Pharmaceuticals